

Black string traveling waves

David Garfinkle

Dept. of Physics, Oakland University, Rochester, MI 48309

Abstract:

A family of solutions to low energy string theory is found. These solutions represent waves traveling along “extremal black strings.”

1. Introduction

The low energy limit of string theory gives a set of classical equations, similar to Einstein's equations, for a metric and other fundamental fields. Several solutions of these equations have been found, including some representing "black strings"¹ that is extended objects with horizons. In this paper we will find some new solutions to low energy string theory. The solutions are found using a generating technique; starting with a known solution the technique produces a new solution representing waves traveling on the old background spacetime. There is no restriction on the amplitude of the waves and, in particular, no approximation in which the waves are considered "small" is used. The background solutions we will use are black strings; so the new solutions represent waves traveling on a black string background.

The fields in the low energy string theory lagrangian are the sigma model metric $\bar{g}_{ab}$, the dilaton ϕ and the axion field strength H_{abc} . Here H_{abc} is derived from a potential B_{ab} by $\mathbf{H} = d\mathbf{B}$. The action in D dimensions is

$$S = \int d^D x \sqrt{\bar{g}} e^{\phi} \left[\bar{R} + \nabla^a \phi \nabla_a \phi - \frac{1}{12} H_{abc} H^{abc} \right] . \quad (1)$$

Here $\bar{R}$ is the scalar curvature of $\bar{g}_{ab}$ and all indicies are raised and lowered with $\bar{g}_{ab}$.

To use the generating technique we will need to write our expressions in terms of the Einstein metric g_{ab} rather than the sigma model metric $\bar{g}_{ab}$. Define the number n , scalar ψ and metric g_{ab} by

$$n \equiv D - 4 , \quad (2)$$

$$\psi \equiv \frac{\phi}{n + 2} , \quad (3)$$

$$g_{ab} \equiv e^{2\psi} \bar{g}_{ab} . \quad (4)$$

In terms of these variables the action is

$$S = \int d^{n+4} x \sqrt{g} \left[R - (n + 2) \nabla^a \psi \nabla_a \psi - \frac{1}{12} e^{4\psi} H_{abc} H^{abc} \right] . \quad (5)$$

Here, and in all subsequent equations, indices are raised and lowered with the Einstein metric g_{ab} . From this action it follows that the equations for the fields are

$$\nabla^a \nabla_a \psi - \frac{1}{6(n+2)} e^{4\psi} H_{abc} H^{abc} = 0 \quad , \quad (6)$$

$$\nabla_a \left(e^{4\psi} H^{abc} \right) = 0 \quad , \quad (7)$$

$$\begin{aligned} R^a_b &= (n+2) \nabla^a \psi \nabla_b \psi + \frac{1}{4} H^{acd} H_{bcd} \\ &\quad - \frac{1}{6(n+2)} \delta^a_b H^{cde} H_{cde} \quad . \end{aligned} \quad (8)$$

We will find solutions of equations (6-8) using the traveling wave generating technique of reference 2 . The technique works as follows: let (g_{ab}, ψ, B_{ab}) be a solution of equations (6-8) that has a null, hypersurface orthogonal Killing vector k^a . Then there is a scalar A such that

$$\nabla_a k_b = k_{[a} \nabla_{b]} \ln A \quad . \quad (9)$$

Define the new metric $\tilde{g}_{ab}$ by

$$\tilde{g}_{ab} \equiv g_{ab} + A \Phi k_a k_b \quad . \quad (10)$$

Let the scalar Φ satisfy the equations

$$k^a \nabla_a \Phi = 0 \quad , \quad (11)$$

$$\nabla^a \nabla_a \Phi = 0 \quad . \quad (12)$$

Our new solution is $(\tilde{g}_{ab}, \psi, B_{ab})$, that is the metric $\tilde{g}_{ab}$ instead of g_{ab} and the same matter fields as in the background solution. It follows from equation (11) that k^a is also a null Killing vector for the new solution. So any disturbances in the new solution must propagate at the speed of light without changing their amplitude or shape. That is, the new solution is a traveling wave.

One can show that $(\tilde{g}_{ab}, \psi, B_{ab})$ satisfies equations (6-8) as follows: a calculation of the Ricci tensor $\tilde{R}^a_b$ of the metric $\tilde{g}_{ab}$ shows² that $\tilde{R}^a_b - R^a_b$ is proportional to $\nabla_a \nabla^a \Phi$. It then follows from equation (12) that $\tilde{R}^a_b = R^a_b$. One then needs to check that raising the indices of H_{abc} with the new metric is the same as in the old metric. It then follows that equation (8) is satisfied. Now the wave operator $\nabla_a \nabla^a$ for scalars invariant under the null translation is the same in the new metric as in the old; so equation (6) is satisfied. Similarly the divergence of an antisymmetric tensor is the same in both metrics; so equation (7) is satisfied.

Therefore all one needs to do, in order to produce a new solution, is solve equations (11) and (12) on the old spacetime. In order to apply this technique the old spacetime must have a null Killing vector. The “extremal black string” solutions of reference 1 have a null Killing vector. In the next section we will apply the generating technique to these solutions.

2. Traveling waves in $D \geq 5$ dimensions

The extremal black string solutions^{1,3} in $D \geq 5$ dimensions are given by the following expressions:

$$\psi = \frac{1}{n+2} \ln \left(1 + \frac{M}{r^n} \right) , \quad (13)$$

$$\mathbf{B} = \left(1 - e^{-(n+2)\psi} \right) du \wedge dv , \quad (14)$$

$$ds^2 = 2e^{-n\psi} dudv + e^{2\psi} \left(dr^2 + r^2 d\Omega_{n+1}^2 \right) . \quad (15)$$

Here $d\Omega_{n+1}^2$ is the metric of the unit $n+1$ sphere and M is a constant.

The null Killing vector is

$$k^a = \left(\frac{\partial}{\partial v} \right)^a \quad (16)$$

and the scalar A is $A = e^{n\psi}$.

We first verify that H_{abc} with the indicies raised by the new metric is the same as with the old metric. The inverse of the new metric is

$$\left(\tilde{g}^{-1}\right)^{ab} = g^{ab} - A\Phi k^a k^b . \quad (17)$$

From equations (14) and (16) it follows that there is an s_a such that $k^a s_a = 0$ and

$$k^a H_{abc} = s_{[b} k_{c]} . \quad (18)$$

It then follows from equation (17) that H_{abc} with its indicies raised by the new metric is the same as H^{abc} .

The new metric is given by

$$d\tilde{s}^2 = ds^2 + e^{-n\psi} \Phi du^2 \quad (19)$$

where Φ satisfies equations (11) and (12). Equation (11) is just

$$\frac{\partial \Phi}{\partial v} = 0 . \quad (20)$$

so Φ depends only on the coordinates u , r and the $n+1$ sphere coordinates. We seek a solution of the form

$$\Phi = f(u) P(r) Y(\theta_i) \quad (21)$$

where the θ_i are the coordinates on the $n+1$ sphere. Equation (12) then becomes

$$0 = e^{-2\psi} f \left[\frac{Y}{r^{n+1}} \frac{d}{dr} \left(r^{n+1} \frac{dP}{dr} \right) + \frac{P}{r^2} D^2 Y \right] . \quad (22)$$

Here D^2 is the Laplacian operator on the $n+1$ sphere. This equation places no restrictions on $f(u)$. Essentially the function f is the profile of the wave and can be chosen arbitrarily. From equation (22) it follows that Y must be an eigenfunction of D^2 . The eigenvalues of D^2 on the $n+1$ sphere are $-l(l+n)$ where l is a nonnegative integer. The function Y is

then Y_l , an $n + 1$ spherical harmonic corresponding to the integer l . It then follows that P satisfies the equation

$$\frac{1}{r^{n+1}} \frac{d}{dr} \left(r^{n+1} \frac{dP}{dr} \right) - \frac{l(l+n)}{r^2} P = 0 \quad . \quad (23)$$

The solutions are $P = r^l$ or $P = r^{-(l+n)}$. The metric is then

$$\begin{aligned} d\tilde{s}^2 = e^{-n\psi} & \left(2 du dv + f(u) r^\beta Y_l(\theta_i) du^2 \right) \\ & + e^{2\psi} \left(dr^2 + r^2 d\Omega_{n+1}^2 \right) \end{aligned} \quad (24)$$

where $\beta = l$ or $\beta = -(l+n)$.

3. Traveling waves in 3 dimensions

The extremal black string in 3 dimensions is given by³

$$\psi = \ln \left(\sqrt{k} r \right) \quad (25)$$

where k is a constant;

$$\mathbf{B} = \frac{M}{r} du \wedge dv \quad , \quad (26)$$

$$ds^2 = 2 k r^2 \left(1 - \frac{M}{r} \right) du dv + \frac{1}{4} k^2 \left(1 - \frac{M}{r} \right)^{-2} dr^2 \quad . \quad (27)$$

Once again the null Killing vector is $k^a = (\partial/\partial v)^a$. The scalar A is now $A = [k^2(r^2 - Mr)]^{-1}$.

As before, the three form H_{abc} is the same with its indicies raised by the new metric as with the old metric. The new metric is

$$d\tilde{s}^2 = ds^2 + r^2 \left(1 - \frac{M}{r} \right) \Phi du^2 \quad . \quad (28)$$

Equation (11) tells us that Φ depends only on r and u . We seek a solution of the form

$$\Phi = f(u) P(r) \quad . \quad (29)$$

Equation (12) then becomes

$$0 = f(u) \frac{d}{dr} \left[(r - M)^2 \frac{dP}{dr} \right] . \quad (30)$$

As before, this equation places no restrictions on $f(u)$. The solutions for $P(r)$ are $P = 1$ or $P = (r - M)^{-1}$. The first solution gives a new metric equivalent to the old metric. So the only physically distinct solution is

$$d\tilde{s}^2 = k r^2 \left(1 - \frac{M}{r} \right) du dv + r f(u) du^2 + \frac{1}{4} k^2 \left(1 - \frac{M}{r} \right)^{-2} dr^2 . \quad (31)$$

4. Properties of the metrics

The extremal black strings are singular at $r = 0$. Therefore the traveling wave solutions are also singular at $r = 0$. One might interpret the extremal black string as the field of a straight fundamental string. In that case the traveling wave solutions would represent propagating modes of the fundamental string.

Some three dimensional black string solutions come from exact conformal field theories.⁴ It would be interesting to see whether there is an exact conformal field theory whose low energy limit is the three dimensional black string traveling wave.

Acknowledgement

I would like to thank Gary Horowitz and Tanmay Vachaspati for helpful discussions.

References

- ¹G. Horowitz and A. Strominger, Nucl. Phys. **B360**, 197 (1991).
- ²D. Garfinkle and T. Vachaspati, Phys. Rev. D **42**, 1960 (1990).
- ³J. Horne, G. Horowitz and A. Steif, Phys. Rev. Lett. **68**, 568 (1992).
- ⁴J. Horne and G. Horowitz, Nucl. Phys. **B368**, 444 (1992).